## Amendments to the Claims:

This listing will replace all prior versions, and listings, of claims in the application:

## **Listing of Claims:**

Claim 1 (Currently amended): A method for directly separating and purifying polyhydroxyalkanoates in cells from <u>a</u> bacterial fermentation liquid, <u>which comprises the steps of:</u> eomprisings:

- (1) <u>a) physically pretreating of the a fermentation liquid containing cells to cause the walls of the cells with physical method</u> to break; cell wall;
- (2) <u>b)</u> adjusting the pH value of the pretreated fermentation liquid from step a) to alkaline; an alkaline condition;
- (3) c) adding anionic surfactant and reacting under to the solution of step b) and subjecting the solution to agitation;
- (4) <u>d)</u> separating and extracting coagulated precipitate from the solution in step c); and reaction liquid;
- (5) e) washing and drying; drying the coagulated precipitate,

Wherein wherein the physical method pretreatment includes mechanical mechanically breaking the cell walls or ultrasonically breaking the cell walls, and wherein steps b) and c) are and ultrasonic breaking; the mechanical breaking includes ball milling or high pressure homogenization; the sequence of adjusting pH and adding surfactant is interchangeable.

Claim 2 (Currently amended): The method according to <u>claim 1</u>, <u>claim 1</u>, wherein <u>a</u> coagulating agent <u>can be</u> is added in the step <u>c</u>). (3).

Claim 3 (Currently amended): The method according to <u>claim 1</u>, <u>claim 1</u>, wherein the <u>physical</u> pretreatment comprises mechanically <u>mechanical</u> breaking the <u>cell walls using is ball milling</u>.

Claim 4 (Currently amended): The method according to <u>claim 1</u>, <u>claim 1</u>, wherein <u>the pH is adjusted</u> in step b) of the pretreated fermentation liquid is adjusted to 8-13.

Claim 5 (Currently amended): The method according to <u>claim 1</u>, <u>claim 1</u> or 4, wherein the <u>alkaline</u> substance used to adjust pH is the pH is adjusted in step b) by adding at least one of NaOH, Na<sub>2</sub>CO<sub>3</sub>, NaHCO<sub>3</sub> solid/aqueous solution or ammonia to the pretreated fermentation liquid. water.

Claim 6 (Currently amended): The method according to <u>claim 1</u>, <u>elaim 1</u>, wherein the anionic surfactant <u>added in step c</u>) is <u>at least one of</u> olefin sulfonate, fatty alcohol sulfate, fatty alcohol polyoxyethylene ether sulfate, fatty alcohol polyoxyethylene ether or alkylphenol polyoxyethylene <u>ether</u>, <u>ether</u>, its quantity is 0.5-20% (W/V) of the fermentation liquid.

Claim 7 (Currently amended): The method according to <u>claim 2</u>, <u>claim 1</u>, wherein the coagulating agent is <u>selected from comprises at least one of sodium polyacrylate</u>, modified starch or <u>polyamine</u>. polyamine, its quantity is 0.5-20% (W/V) of the fermentation liquid.

Claim 8 (Currently amended): The method according to <u>claim 1</u>, <u>claim 1</u>, wherein <u>in step c</u>) the reaction temperature <u>of the solution is adjusted to 10 - 70°C</u>. <u>under agitation is 10 - 70°C</u>, the time is <u>5 60min</u>.

Claim 9 (Currently amended): The method according to <u>claim 1</u>, <u>elaim 1</u>, wherein <u>in step d</u>) the <u>method for separating and extracting</u> coagulated precipitate <u>is separated and extracted by means of a from reaction liquid is selected from centrifuge</u>, filter press or vacuum suction filtration.

Claim 10 (New): The method according to claim 1, claim1, wherein the physical pretreatment comprises high pressure homogenization.

Claim 11 (New): The method according to claim 4, wherein the pH is adjusted in step b) by adding at least one of NaOH, Na<sub>2</sub>CO<sub>3</sub>, NaHCO<sub>3</sub> or ammonia to the pretreated fermentation liquid.

Claim 12 (New): The method according to claim 6, wherein the amount of anionic surfactant added in step c) is 0.5-20% (W/V) of the fermentation liquid.

Claim 13 (New): The method according to claim 7, wherein the amount of coagulating agent added is 0.5-20% (W/V) of the fermentation liquid.

Claim 14 (New): The method according to claim 1, wherein in step c) the solution is subjected to agitation for 5 - 60 min.

Claim 15 (New): The method according to claim 8, wherein in step c) the solution is subjected to agitation for 5 - 60 min.

Claim 16 (New): The method according to claim 5, wherein the NaOH, Na<sub>2</sub>CO<sub>3</sub>, NaHCO<sub>3</sub> is added as a solid.

Claim 17 (New): The method according to claim 5, wherein the NaOH, Na<sub>2</sub>CO<sub>3</sub>, NaHCO<sub>3</sub> is added as a solution.

Claim 18 (New): The method according to claim 5, wherein the ammonia comprises ammonia water.